

IL3RA rabbit monoclonal antibody

Catalog # H00003563-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human IL3RA peptide using ARM Technology.
Immunogen	A synthetic peptide of human IL3RA is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human IL3RA peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — IL3RA

Entrez GeneID	3563
GeneBank Accession#	IL3RA
Gene Name	IL3RA
Gene Alias	CD123, IL3R, IL3RAY, IL3RX, IL3RY, MGC34174, hIL-3Ra
Gene Description	interleukin 3 receptor, alpha (low affinity)
Omim ID	308385 430000
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudoautosomal region on chromosomes X or Y. [provided by RefSeq]
Other Designations	CD123 antigen Interleukin-3 receptor OTTHUMP00000022828 interleukin 3 receptor, alpha

Pathway

- [Apoptosis](#)
- [Cytokine-cytokine receptor interaction](#)
- [Hematopoietic cell lineage](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Chorioamnionitis](#)
- [Fetal Membranes](#)

- [Genetic Predisposition to Disease](#)
- [Obstetric Labor](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Schizophrenia](#)
- [Thyroid Neoplasms](#)