


Full-Length

IL15 (Human) Recombinant Protein (P02)

Catalog # H00003600-P02

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human IL15 full-length ORF (NP_000576.1, 1 a.a. - 162 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MRISKPHLRISIQCYLCLLNSHFLTEAGIHVFILGCF SAGLPKTEANWVNVISDLKKIEDLIQSMHID ATLYTESDVHPSCKVTAMKCFLEELQVISLES GDASIHDTVENLILANNSLSSNGNVTESGCKECE ELEEKNIKEFLQSFVHIVQMFINTS
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	44.5
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — IL15

Entrez GeneID [3600](#)

GeneBank Accession# [NM_000585.2](#)

Protein Accession# [NP_000576.1](#)

Gene Name IL15

Gene Alias IL-15, MGC9721

Gene Description interleukin 15

Omim ID [600554](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported. [provided by RefSeq]

Other Designations OTTHUMP00000164617

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Arthritis](#)
- [Asthma](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Connective Tissue Diseases](#)
- [Coronary Disease](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Diseases](#)
- [Fractures](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Hypersensitivity](#)
- [Hypertrophy](#)
- [Infant](#)
- [Infection](#)

- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasm](#)
- [Obesity](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Psoriasis](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Skin Diseases](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)