

DNAxPAb



## KCNA4 DNAxPab

Catalog # H00003739-W01P Size 200 ug

Specification	
Product Description	Rabbit polyclonal antibody raised against a partial-length human KCNA4 DNA using DNAx™ Immun e technology.
Technology	DNAx™ Immune
Immunogen	Extracellular membrane domain (ECD) human DNA
Host	Rabbit
Reactivity	Human
Purification	Protein A
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot (Transfected lysate)
  <u>Protocol Download</u>
- Immunofluorescence (Transfected cell)
- Flow Cytometry (Transfected cell)

## Gene Info — KCNA4

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😭 Abnova	Product Information
Entrez GenelD	3739
GeneBank Accession#	<u>NM_002233.2</u>
Protein Accession#	<u>NP_002224.1</u>
Gene Name	KCNA4
Gene Alias	HBK4, HK1, HPCN2, HUKII, KCNA4L, KCNA8, KV1.4, PCN2
Gene Description	potassium voltage-gated channel, shaker-related subfamily, member 4
Omim ID	<u>176266</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Potassium channels represent the most complex class of voltage-gated ion channels from both fu nctional and structural standpoints. Their diverse functions include regulating neurotransmitter rele ase, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth mus cle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in Drosophila, and each has been shown to have human ho molog(s). This gene encodes a member of the potassium channel, voltage-gated, shaker-related subfamily. This member contains six membrane-spanning domains with a shaker-type repeat in th e fourth segment. It belongs to the A-type potassium current class, the members of which may be i mportant in the regulation of the fast repolarizing phase of action potentials in heart and thus may i nflunce the duration of cardiac action potential. The coding region of this gene is intronless, and th e gene is clustered with genes KCNA3 and KCNA10 on chromosome 1. [provided by RefSeq
Other Designations	cardiac potassium channel fetal skeletal muscle potassium channel potassium channel 2 potassiu m voltage-gated channel, shaker-related subfamily, member 4-like rapidly inactivating potassium channel shaker-related potassium channel Kv1.4 type A potassium c