


LIMK1 monoclonal antibody (M01), clone 1A8

Catalog # H00003984-M01


Size 100 ug

Applications




Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in PC-12 (Cat # L012V1).




Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in NIH/3T3 (Cat # L018V1).




Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in SW-13 (Cat # L005V1).




Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8 Western Blot analysis of LIMK1 expression in HepG2 (Cat # L019V1).




Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in Raw 264.7 (Cat # L024V1).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunoperoxidase of monoclonal antibody to LIMK1 on formalin-fixed paraffin-embedded human stomach. [antibody concentration 3 ug/ml]



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged LIMK1 is approximately 0.03ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.74 KDa).

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant LIMK1.

Immunogen

LIMK1 (NP_002305, 548 a.a. ~ 647 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence	ADPDYLPRTMDFGLNVRGFLDRYCPNCPSPSFFPITVRCCDLDPKRPSPFVKLEHWLETLRMHL AGHLPLGPQLEQLDRGFWEYRRGESGLPAHPEVPD
Host	Mouse
Reactivity	Human, Mouse, Rat
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in PC-12(Cat # L012V1).

[Protocol Download](#)

- Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in NIH/3T3 (Cat # L018V1).

[Protocol Download](#)

- Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in SW-13 (Cat # L005V1).

[Protocol Download](#)

- Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8 Western Blot analysis of LIMK1 expression in HepG2 (Cat # L019V1).

[Protocol Download](#)

- Western Blot (Cell lysate)

LIMK1 monoclonal antibody (M01), clone 1A8. Western Blot analysis of LIMK1 expression in Raw 264.7(Cat # L024V1).

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunoperoxidase of monoclonal antibody to LIMK1 on formalin-fixed paraffin-embedded human stomach. [antibody concentration 3 ug/ml]

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged LIMK1 is approximately 0.03ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — LIMK1

Entrez GeneID [3984](#)

GeneBank Accession# [NM_002314](#)

Protein Accession# [NP_002305](#)

Gene Name LIMK1

Gene Alias LIMK

Gene Description LIM domain kinase 1

Omim ID [601329](#)

Gene Ontology [Hyperlink](#)

Gene Summary

There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is likely to be a component of an intracellular signaling pathway and may be involved in brain development. LIMK1 hemizygosity is implicated in the impaired visuospatial constructive cognition of Williams syndrome. [provided by RefSeq]

Other Designations LIM motif-containing protein kinase|OTTHUMP00000025066

Pathway

- [Axon guidance](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Brain Ischemia](#)
- [Cerebral Hemorrhage](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Genetic Predisposition to Disease](#)
- [Intracranial Aneurysm](#)
- [Metabolic Syndrome X](#)
- [Renal Insufficiency](#)
- [Stroke](#)
- [Werner syndrome](#)