

MARCKS rabbit monoclonal antibody

Catalog # H00004082-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human MARCKS peptide using ARM Technology.
Immunogen	A synthetic peptide of human MARCKS is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human MARCKS peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — MARCKS	
Entrez GenelD	4082
GeneBank Accession#	MARCKS
Gene Name	MARCKS
Gene Alias	80K-L, FLJ14368, FLJ90045, MACS, PKCSL, PRKCSL
Gene Description	myristoylated alanine-rich protein kinase C substrate
Omim ID	<u>177061</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a substrate for protein kinase C. It is localized to the plasma membrane and is an actin filament crosslinking protein. Phosphorylation by protein kinase C or bi nding to calcium-calmodulin inhibits its association with actin and with the plasma membrane, lea ding to its presence in the cytoplasm. The protein is thought to be involved in cell motility, phagocy tosis, membrane trafficking and mitogenesis. [provided by RefSeq
Other Designations	OTTHUMP00000017045 myristoylated alanine-rich protein kinase C substrate (MARCKS, 80K-L) phosphomyristin

Pathway

• Fc gamma R-mediated phagocytosis