


MDH2 monoclonal antibody (M01), clone 1G12

Catalog # H00004191-M01


Size 100 ug

Applications




Western Blot (Cell lysate)

MDH2 monoclonal antibody (M01), clone 1G12. Western Blot analysis of MDH2 expression in IMR-32 (Cat # L008V1).



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged MDH2 is 0.3 ng/ml as a capture antibody.



Western Blot detection against Immunogen (38.17 KDa).

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant MDH2.

Immunogen	MDH2 (NP_005909, 134 a.a. ~ 246 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	EAMICVIANPVNSTIPITAEVFKKHGVYNPNKIFGVTTLDIVRANTFVAELKGLDPARVNVVPVIGGHA GKTIPLISQCTPKVDFPQDQLTALTGRIQEAGTEVVKAKAGAGS
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (96); Rat (96)
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (38.17 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

MDH2 monoclonal antibody (M01), clone 1G12. Western Blot analysis of MDH2 expression in IMR-32(Cat # L008V1).

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged MDH2 is 0.3 ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — MDH2

Entrez GeneID [4191](#)

GeneBank Accession#	NM_005918
Protein Accession#	NP_005909
Gene Name	MDH2
Gene Alias	M-MDH, MDH, MGC:3559, MOR1
Gene Description	malate dehydrogenase 2, NAD (mitochondrial)
Omim ID	154100
Gene Ontology	Hyperlink
Gene Summary	Malate dehydrogenase catalyzes the reversible oxidation of malate to oxaloacetate, utilizing the NAD/NADH cofactor system in the citric acid cycle. The protein encoded by this gene is localized to the mitochondria and may play pivotal roles in the malate-aspartate shuttle that operates in the metabolic coordination between cytosol and mitochondria. [provided by RefSeq]
Other Designations	mitochondrial malate dehydrogenase

Pathway

- [Biosynthesis of alkaloids derived from histidine and purine](#)
- [Biosynthesis of alkaloids derived from ornithine](#)
- [Biosynthesis of alkaloids derived from shikimate pathway](#)
- [Biosynthesis of alkaloids derived from terpenoid and polyketide](#)
- [Biosynthesis of phenylpropanoids](#)
- [Biosynthesis of plant hormones](#)
- [Biosynthesis of terpenoids and steroids](#)
- [Carbon fixation in photosynthetic organisms](#)
- [Citrate cycle \(TCA cycle\)](#)
- [Glyoxylate and dicarboxylate metabolism](#)
- [Metabolic pathways](#)
- [Pyruvate metabolism](#)
- [Reductive carboxylate cycle \(CO2 fixation\)](#)

Disease

- [Tobacco Use Disorder](#)