


MDM4 (Human) IP-WB Antibody Pair

Catalog # H00004194-PW1 Size 1 Set

Applications



Immunoprecipitation of MDM4 transfected lysate using rabbit polyclonal anti-MDM4 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-MDM4.

Specification	
Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Interspecies Antigen Sequence	Mouse (83%); Rat (83%)
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of MDM4 transfected lysate using rabbit polyclonal anti-MDM4 and Protein A M agnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-MDM4.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-MDM4 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-MDM4 (50 ug)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use.

Applications



Immunoprecipitation-Western Blot

Protocol Download

Gene Info — MDM4	
Entrez GenelD	4194
Gene Name	MDM4
Gene Alias	DKFZp781B1423, HDMX, MDMX, MGC132766, MRP1
Gene Description	Mdm4 p53 binding protein homolog (mouse)
Omim ID	602704
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The human MDM4 gene, which plays a role in apoptosis, encodes a 490-amino acid protein cont aining a RING finger domain and a putative nuclear localization signal. The MDM4 putative nuclear localization signal, which all Mdm proteins contain, is located in the C-terminal region of the prot ein. The mRNA is expressed at a high level in thymus and at lower levels in all other tissues tested . MDM4 protein produced by in vitro translation interacts with p53 via a binding domain located in the N-terminal region of the MDM4 protein. MDM4 shows significant structural similarity to p53-binding protein MDM2. Two transcript variants, one protein-coding and the other likely not to be protein-coding, have been found for this gene. [provided by RefSeq
Other Designations	MDM4-related protein 1 Mdm4, transformed 3T3 cell double minute 4, p53 binding protein double minute 4, human homolog of; p53-binding protein mouse double minute 4 homolog p53-binding protein

Pathway

• p53 signaling pathway

Disease

- Breast cancer
- Breast Neoplasms
- Carcinoma
- Genetic Predisposition to Disease



- Head and Neck Neoplasms
- Narcolepsy
- Neoplasm Recurrence
- Neoplasms
- Tobacco Use Disorder