


MaxPab®

MMP7 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00004316-B01P

Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of MMP7 expression in transfected 293T cell line ([H00004316-T01](#)) by MMP7 MaxPab polyclonal antibody.

Lane 1: MMP7 transfected lysate(29.37 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Mouse polyclonal antibody raised against a full-length human MMP7 protein.
Immunogen	MMP7 (NP_002414.1, 1 a.a. ~ 267 a.a) full-length human protein.
Sequence	MRLTVLCVAVCLLPGLSLALPLPQEAGGMSELQWEQAQDYLRFYLYDSETKNANSLEAKLKEMQK FFGLPITGMLNSRVIEIMQKPRCGVPDVAEYSLFPNSPKWTSKVVTYRVSYTRDLPHITVDRLVSK ALNMWGKEIPLHFRKVVGWGTADIMIGFARGAHGDSYPFDGPGNTLAHAFAPGTGLGGDAHFD ERWTDGSSSLGINFLYAAATHELGHSLGMGHSSDPNAVVMYPTYGNGDPQNFKLSQDDIKGIQKLYGK RSNSRKK
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (70); Rat (70)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of MMP7 expression in transfected 293T cell line ([H00004316-T01](#)) by MMP7 MaxPab polyclonal antibody.

Lane 1: MMP7 transfected lysate(29.37 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — MMP7

Entrez GeneID [4316](#)

GeneBank Accession# [NM_002423](#)

Protein Accession# [NP_002414.1](#)

Gene Name MMP7

Gene Alias MMP-7, MPST1, PUMP-1

Gene Description matrix metalloproteinase 7 (matrilysin, uterine)

Omim ID [178990](#)

Gene Ontology [Hyperlink](#)

Gene Summary Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans, fibronectin, elastin and casein and differs from most MMP family members in that it lacks a conserved C-terminal protein domain. The enzyme is involved in wound healing, and studies in mice suggest that it regulates the activity of defensins in intestinal mucosa. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq]

Other Designations matrin|matrix metalloproteinase 7|matrix metalloproteinase 7 (matrilysin, uterine)|uterine matrilysin

Pathway

- [Wnt signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Adenomatous Polyps](#)
- [Arthritis](#)
- [Astrocytoma](#)
- [Atherosclerosis](#)
- [Atrophy](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis Obliterans](#)
- [Carcinoma](#)
- [Chorioamnionitis](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Disease Progression](#)
- [Duodenal Ulcer](#)
- [Endometrial Neoplasms](#)
- [Endometriosis](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)
- [Gastritis](#)
- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)

- [Heart Neoplasms](#)
- [Helicobacter Infections](#)
- [Hepatitis C](#)
- [Hypercholesterolemia](#)
- [Liver Cirrhosis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Sclerosis](#)
- [Nasopharyngeal Neoplasms](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Precancerous Conditions](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Stomach Ulcer](#)
- [Thyroid Neoplasms](#)

- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)