
MYOG monoclonal antibody (M01), clone 2B7

Catalog # H00004656-M01 Size 100 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of MYOG expression in transfected 293T cell line by MYOG monoclonal antibody (M01), clone 2B7.

Lane 1: MYOG transfected lysate(25 KDa). Lane 2: Non-transfected lysate.

Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged MYOG is approximately 0.03ng/ml as a capture antibody.




RNAi Knockdown (Antibody validated)

Western blot analysis of MYOG over-expressed 293 cell line, cotransfected with MYOG Validated Chimera RNAi (Cat # H00004656-R01V) (Lane 2) or non-transfected control (Lane 1). Blot probed with MYOG monoclonal antibody (M01), clone 2B7 (Cat # H00004656-M01). GAPDH (36.1 kDa) used as specificity and loading control.

😵 Abnova

Product Information



Western Blot detection against Immunogen (50.38 KDa).

Specification	
Product Description	Mouse monoclonal antibody raised against a full length recombinant MYOG.
Immunogen	MYOG (AAH53899, 1 a.a. ~ 224 a.a) full-length recombinant protein with GST tag. MW of the GST ta g alone is 26 KDa.
Sequence	MELYETSPYFYQEPRFYDGENYLPVHLQGFEPPGYERTELTLSPEAPGPLEDKGLGTPEHCPGQ CLPWACKVCKRKSVSVDRRRAATLREKRRLKKVNEAFEALKRSTLLNPNQRLPKVEILRSAIQYIE RLQALLSSLNQEERDLRYRGGGGPQPGVPSECSSHSASCSPEWGSALEFSANPGDHLLTADPT DAHNLHSLTSIVDSITVEDVSVAFPDETMPN
Host	Mouse
Reactivity	Human
lsotype	lgG1 kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (50.38 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

• Western Blot (Transfected lysate)

Western Blot analysis of MYOG expression in transfected 293T cell line by MYOG monoclonal antibody (M01), clone 2B7.

Lane 1: MYOG transfected lysate(25 KDa). Lane 2: Non-transfected lysate.

Protocol Download



- Western Blot (Recombinant protein)
 Protocol Download
- Sandwich ELISA (Recombinant protein)
 Detection limit for recombinant GST tagged MYOG is approximately 0.03ng/ml as a capture antibody.
 <u>Protocol Download</u>
- ELISA
- RNAi Knockdown (Antibody validated)

Western blot analysis of MYOG over-expressed 293 cell line, cotransfected with MYOG Validated Chimera RNAi (Cat # H00004656-R01V) (Lane 2) or non-transfected control (Lane 1). Blot probed with MYOG monoclonal antibody (M01), clone 2B7 (Cat # H00004656-M01). GAPDH (36.1 kDa) used as specificity and loading control.

Protocol Download

Gene Info — MYOG

Entrez GenelD	<u>4656</u>
GeneBank Accession#	<u>BC053899</u>
Protein Accession#	AAH53899
Gene Name	MYOG
Gene Alias	MYF4, MYOGENIN, bHLHc3
Gene Description	myogenin (myogenic factor 4)
Omim ID	<u>159980</u>
Gene Ontology	Hyperlink
Gene Summary	Myogenin is a muscle-specific transcription factor that can induce myogenesis in a variety of cell t ypes in tissue culture. It is a member of a large family of proteins related by sequence homology, t he helix-loop-helix (HLH) proteins. It is essential for the development of functional skeletal muscle. [provided by RefSeq
Other Designations	Myogenic factor-4; myogenin OTTHUMP00000039094 myogenic factor 4 myogenin