


CEACAM6 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00004680-T01


Size 100 uL

Applications



SDS-PAGE Gel

CEACAM6 transfected lysate.



Western Blot

Lane 1: CEACAM6 transfected lysate (37.2 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-CEACAM6 full-length

Host Human

Theoretical MW (kDa) 37.2

Quality Control Testing Transient overexpression cell lysate was tested with Anti-CEACAM6 antibody ([H00004680-B01](#)) by Western Blots.
SDS-PAGE Gel
CEACAM6 transfected lysate.
Western Blot
Lane 1: CEACAM6 transfected lysate (37.2 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — CEACAM6

Entrez GeneID	4680
GeneBank Accession#	BC005008.1
Protein Accession#	-
Gene Name	CEACAM6
Gene Alias	CD66c, CEAL, NCA
Gene Description	carcinoembryonic antigen-related cell adhesion molecule 6 (non-specific cross reacting antigen)
Omim ID	163980
Gene Ontology	Hyperlink
Gene Summary	Carcinoembryonic antigen (CEA; MIM 114890) is one of the most widely used tumor markers in serum immunoassay determinations of carcinoma. An apparent lack of absolute cancer specificity for CEA probably results in part from the presence in normal and neoplastic tissues of antigens that share antigenic determinants with the 180-kD form of CEA (Barnett et al., 1988 [PubMed 3220478]). For background information on the CEA family of genes, see CEACAM1 (MIM 109770). [supplied by OMIM]
Other Designations	-

Disease

- [Genetic Predisposition to Disease](#)
- [Meningococcal Infections](#)