


NFIC 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00004782-T01


Size 100 uL

Applications



SDS-PAGE Gel

NFIC transfected lysate.



Western Blot

Lane 1: NFIC transfected lysate (47.19 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-NFIC full-length

Host Human

Theoretical MW (kDa) 47.19

Quality Control Testing Transient overexpression cell lysate was tested with Anti-NFIC antibody ([H00004782-B01](#)) by Western Blots.
 SDS-PAGE Gel
 NFIC transfected lysate.
 Western Blot
 Lane 1: NFIC transfected lysate (47.19 KDa)
 Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — NFIC

Entrez GeneID[4782](#)**GeneBank Accession#**[NM_005597.2](#)**Protein Accession#**[NP_005588.2](#)**Gene Name**

NFIC

Gene Alias

CTF, CTF5, MGC20153, NF-I, NFI

Gene Description

nuclear factor I/C (CCAAT-binding transcription factor)

Omim ID[600729](#)**Gene Ontology**[Hyperlink](#)**Other Designations**

CCAAT-binding transcription factor|nuclear factor I/C

Disease

- [Tobacco Use Disorder](#)