


PAK3 monoclonal antibody (M07), clone 1H7

Catalog # H00005063-M07


Size 100 ug

Applications



Western Blot (Cell lysate)

PAK3 monoclonal antibody (M07), clone 1H7 Western Blot analysis of PAK3 expression in NIH/3T3 (Cat # L018V1).



Immunofluorescence

Immunofluorescence of monoclonal antibody to PAK3 on NIH/3T3 cell. [antibody concentration 10 ug/ml]



Western Blot detection against Immunogen (35.53 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant PAK3.

Immunogen	PAK3 (NP_002569, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	MSDGLDNEEKPPAPPLRMNSNNRDSSALNHSSKPLPMAPEEKNKKARLRSIFPGGGDKTNKKK EKERPEISLPDFEHTIHVGFDVAVTGE
Host	Mouse
Reactivity	Human, Mouse
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.53 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

PAK3 monoclonal antibody (M07), clone 1H7 Western Blot analysis of PAK3 expression in NIH/3T3 (Cat # L018V1).

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

- Immunofluorescence

Immunofluorescence of monoclonal antibody to PAK3 on NIH/3T3 cell. [antibody concentration 10 ug/ml]

Gene Info — PAK3

Entrez GeneID	5063
GeneBank Accession#	NM_002578
Protein Accession#	NP_002569
Gene Name	PAK3

Gene Alias	CDKN1A, MRX30, MRX47, OPHN3, PAK3beta, bPAK, hPAK3
Gene Description	p21 protein (Cdc42/Rac)-activated kinase 3
Omim ID	300142 300558
Gene Ontology	Hyperlink
Gene Summary	PAK proteins are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. PAK proteins, a family of serine/threonine p21-activating kinases, serve as targets for the small GTP binding proteins Cdc42 and RAC and have been implicated in a wide range of biological activities. The protein encoded by this gene forms an activated complex with GTP-bound RAS-like (P21), CDC2 and RAC1 proteins which then catalyzes a variety of targets. This protein may be necessary for dendritic development and for the rapid cytoskeletal reorganization in dendritic spines associated with synaptic plasticity. Defects in this gene are the cause of non-syndromic mental retardation X-linked type 30 (MRX30), also called X-linked mental retardation type 47 (MRX47). Alternatively spliced transcript variants encoding different isoforms have been identified . [provided by RefSeq]
Other Designations	OTTHUMP00000023855 OTTHUMP00000062894 beta-PAK oligophrenin-3 p21 (CDKN1A)-activated kinase 3 p21-activated kinase 3 p21-activated kinase-3 serine/threonine-protein kinase PAK 3

Publication Reference

- [A 14-3-3γ dimer-based scaffold bridges CtBP1-S/BARS to PI\(4\)KIIIβ to regulate post-Golgi carrier formation.](#)

Valente C, Turacchio G, Mariggio S, Pagliuso A, Gaibisso R, Di Tullio G, Santoro M, Formiggini F, Spanò S, Piccini D, Polishchuk RS, Colanzi A, Luini A, Corda D.

Nature Cell Biology 2012 Feb; 14(4):343.

Application: IF, Monkey, COS7 cells

Pathway

- [Axon guidance](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Regulation of actin cytoskeleton](#)
- [Renal cell carcinoma](#)
- [T cell receptor signaling pathway](#)

Disease

- [Chronic Disease](#)
- [Cognition Disorders](#)
- [Endometrial Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Neuropsychological Tests](#)
- [Schizophrenia](#)