

PARN rabbit monoclonal antibody

Catalog # H00005073-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human PARN peptide using ARM Technology.
Immunogen	A synthetic peptide of human PARN is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human PARN peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — PARN	
Entrez GenelD	<u>5073</u>
GeneBank Accession#	PARN
Gene Name	PARN
Gene Alias	DAN
Gene Description	poly(A)-specific ribonuclease (deadenylation nuclease)
Omim ID	604212
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a 3'-exoribonuclease, with similarity to the RNase D family of 3'-exonucleases. It prefers poly(A) as the substrate, hence, efficiently degrades poly(A) tails of mR NAs. Exonucleolytic degradation of the poly(A) tail is often the first step in the decay of eukaryotic mRNAs. This protein is also involved in silencing of certain maternal mRNAs during oocyte matur ation and early embryonic development, as well as in nonsense-mediated decay (NMD) of mRNAs that contain premature stop codons. Alternatively spliced transcript variants encoding different is oforms have been found for this gene. [provided by RefSeq
Other Designations	deadenylating nuclease

Pathway

RNA degradation

Disease

Tobacco Use Disorder