


PCNA monoclonal antibody (M06), clone 2A6

Catalog # H00005111-M06


Size 100 ug

Applications




Western Blot (Tissue lysate)

PCNA monoclonal antibody (M06), clone 2A6. Western Blot analysis of PCNA expression in human placenta.




Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PCNA is approximately 0.3ng/ml as a capture antibody.



Immunofluorescence

Immunofluorescence of monoclonal antibody to PCNA on HeLa cell . [antibody concentration 10 ug/ml]



Western Blot detection against Immunogen (37 KDa) .

Specification

Product Description	Mouse monoclonal antibody raised against a full length recombinant PCNA.
Immunogen	PCNA (NP_002583, 78 a.a. ~ 177 a.a) full length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	ILKCAGNEDIITLRAEDNADTLALVFEAPNQEKVSDYEMKLMDL DVEQLGIPEQEYSCVVKMPSGE FARICRDL SHIGDAVVISCAKDGVKFSASGELGN
Host	Mouse
Reactivity	Human
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Tissue lysate)

PCNA monoclonal antibody (M06), clone 2A6. Western Blot analysis of PCNA expression in human placenta.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PCNA is approximately 0.3ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA
- Immunofluorescence

Immunofluorescence of monoclonal antibody to PCNA on HeLa cell . [antibody concentration 10 ug/ml]

Gene Info — PCNA

Entrez GeneID [5111](#)

GeneBank Accession# [NM_002592](#)

Protein Accession# [NP_002583](#)

Gene Name PCNA

Gene Alias MGC8367

Gene Description proliferating cell nuclear antigen

Omim ID [176740](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome. [provided by RefSeq]

Other Designations DNA polymerase delta auxiliary protein|OTTHUMP00000030189|OTTHUMP00000030190|cyclin

Pathway

- [Base excision repair](#)
- [Cell cycle](#)
- [DNA replication](#)

- [Mismatch repair](#)
- [Nucleotide excision repair](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia telangiectasia](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [DNA Damage](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Graft vs Host Disease](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Laryngeal Neoplasms](#)
- [Lung Neoplasms](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neuroma](#)

- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Pharyngeal Neoplasms](#)
- [Pulmonary Disease](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)