


PFDN4 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00005203-T05 Size 100 uL

Applications



SDS-PAGE Gel

PFDN4 transfected lysate

Western Blot

Lane 1: PFDN4 transfected lysate (14.85 KDa). Lane 2: Non-transfected lysate.

| Specification | |
|----------------------------------|------------------------|
| Transfected Cell Line | 293T |
| Plasmid | pCMV-PFDN4 full-length |
| Host | Human |
| Theoretical MW (kDa) | 14.85 |
| Interspecies Antigen Sequence | Mouse (97); Rat (94) |



Product Information

| Quality Control Testing | Transient overexpression cell lysate was tested with Anti-PFDN4 antibody (H00005203-B03) by We stern Blots. SDS-PAGE Gel PFDN4 transfected lysate Western Blot Lane 1: PFDN4 transfected lysate (14.85 KDa). Lane 2: Non-transfected lysate. |
|-------------------------|--|
| Storage Buffer | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue) |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |

Applications

• Western Blot

Gene Info — PFDN4

| Entrez GenelD | <u>5203</u> |
|---------------------|--|
| GeneBank Accession# | <u>BC010953</u> |
| Protein Accession# | AAH10953 |
| Gene Name | PFDN4 |
| Gene Alias | C1, PFD4 |
| Gene Description | prefoldin subunit 4 |
| Omim ID | 604898 |
| Gene Ontology | Hyperlink |
| Gene Summary | This gene encodes a member of the prefoldin beta subunit family. The encoded protein is one of six subunits of prefoldin, a molecular chaperone complex that binds and stabilizes newly synthesiz ed polypeptides, thereby allowing them to fold correctly. The complex, consisting of two alpha and four beta subunits, forms a double beta barrel assembly with six protruding coiled-coils. [provided by RefSeq |
| Other Designations | OTTHUMP0000031316 prefoldin 4 |