PGAM2 rabbit monoclonal antibody

Catalog # H00005224-K

Specification

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human PGAM2 peptide using ARM Technology.
Immunogen	A synthetic peptide of human PGAM2 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human PGAM2 peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download



• ELISA

Gene Info — PGAM2	
Entrez GenelD	<u>5224</u>
GeneBank Accession#	PGAM2
Gene Name	PGAM2
Gene Alias	MGC88743, PGAM-M, PGAMM
Gene Description	phosphoglycerate mutase 2 (muscle)
Omim ID	261670
Gene Ontology	Hyperlink
Gene Summary	Phosphoglycerate mutase (PGAM) catalyzes the reversible reaction of 3-phosphoglycerate (3-PG A) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The PGAM is a dimeric enzyme cont aining, in different tissues, different proportions of a slow-migrating muscle (MM) isozyme, a fast-migrating brain (BB) isozyme, and a hybrid form (MB). This gene encodes muscle-specific PGAM subunit. Mutations in this gene cause muscle phosphoglycerate mutase eficiency, also known as glycogen storage disease X. [provided by RefSeq
Other Designations	Phosphoglycerate mutase, muscle form

Pathway

- Biosynthesis of alkaloids derived from histidine and purine
- Biosynthesis of alkaloids derived from ornithine
- Biosynthesis of alkaloids derived from shikimate pathway
- Biosynthesis of alkaloids derived from terpenoid and polyketide
- Biosynthesis of phenylpropanoids
- Biosynthesis of plant hormones
- Biosynthesis of terpenoids and steroids
- <u>Glycolysis / Gluconeogenesis</u>
- Metabolic pathways