


PLAU monoclonal antibody (M01), clone 3B8

Catalog # H00005328-M01


Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PLAU is approximately 0.1ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.63 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant PLAU.

Immunogen

PLAU (AAH13575, 16 a.a. ~ 115 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence

SDSKGSNELHQVPSNCDCLNGGTCVSNKYFSNIHWCNCPKKFGGQHCEIDKSKTCYEGNGHFY
RGKASTDTMGRPCLPWNSATVLQQTYHAHRSDALQLG

Host

Mouse

Reactivity

Human

Interspecies Antigen Sequence	Mouse (69); Rat (71)
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PLAUI is approximately 0.1ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — PLAUI

Entrez GeneID	5328
GeneBank Accession#	BC013575
Protein Accession#	AAH13575
Gene Name	PLAUI
Gene Alias	ATF, UPA, URK, UROKINASE, u-PA
Gene Description	plasminogen activator, urokinase
Omim ID	104300 191840
Gene Ontology	Hyperlink

Gene Summary

This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active, carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA). HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

OTTHUMP00000019849|U-plasminogen activator|antagonist of uPA|plasminogen activator, urinary|urokinase plasminogen activator|urokinase-type plasminogen activator amino-terminal fragment

Pathway

- [Complement and coagulation cascades](#)

Disease

- [Alveolar Bone Loss](#)
- [Alzheimer disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Bacteremia](#)
- [Blood Platelet Disorders](#)
- [Bronchopulmonary Dysplasia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Chronic Disease](#)
- [Cognition](#)
- [Colorectal Neoplasms](#)

- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Gram-Negative Bacterial Infections](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Insulin Resistance](#)
- [Kidney Calculi](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Metabolic Syndrome X](#)
- [Mitral Valve Prolapse](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Myocardial Infarction](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pneumoconiosis](#)
- [Polycystic Ovary Syndrome](#)

- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Recurrence](#)
- [Respiratory Distress Syndrome](#)
- [Small Cell Lung Carcinoma](#)
- [Thyroid Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Urinary Calculi](#)