


PPARBP monoclonal antibody (M03), clone 1G4

Catalog # H00005469-M03


Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PPARBP is approximately 0.1ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.74 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant PPARBP.

Immunogen

PPARBP (NP_004765, 1391 a.a. ~ 1490 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence

KIIISKHDGGSPSIKAKVTLQKPGESSGEGLRPQMASSKNYGSPLISGSTPKHERGSPSHSKSPAY
TPQNLDSESESGSSIAEKSYQNSPSSDDGIRPLP

Host

Mouse

Reactivity

Human

Interspecies Antigen Sequence	Mouse (94); Rat (94)
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged PPARBP is approximately 0.1ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — MED1

Entrez GeneID	5469
GeneBank Accession#	NM_004774
Protein Accession#	NP_004765
Gene Name	MED1
Gene Alias	CRSP1, CRSP200, DRIP205, DRIP230, MGC71488, PBP, PPARBP, PPARGBP, RB18A, TRA P220, TRIP2
Gene Description	mediator complex subunit 1
Omim ID	604311
Gene Ontology	Hyperlink

Gene Summary

The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. It also regulates p53-dependent apoptosis and it is essential for adipogenesis. This protein is known to have the ability to self-oligomerize. [provided by RefSeq]

Other Designations

PPAR binding protein|PPARG binding protein|thyroid hormone receptor-associated protein complex component TRAP220|thyroid receptor interacting protein 2|vitamin D receptor-interacting protein 230 kD|vitamin D receptor-interacting protein complex component

Disease

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)