


MAP2K2 (Human) Matched Antibody Pair

Catalog # H00005605-AP21 Size 1 Set

Applications



Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.

Specification

Product Description	This antibody pair set comes with a matched antibody pair to detect and quantify the protein level of human MAP2K2.
Reactivity	Human
Quality Control Testing	Standard curve using recombinant protein (H00005605-P01) as an analyte. Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.
Supplied Product	Antibody pair set content: 1. Capture antibody: rabbit MaxPab® affinity purified polyclonal anti-MAP2K2 (100 ug) 2. Detection antibody: mouse purified polyclonal anti-MAP2K2 (20 ug) *Reagents are sufficient for at least 1-2 x 96 well plates using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- ELISA Pair (Recombinant protein)

[Protocol Download](#)

Gene Info — MAP2K2

Entrez GeneID [5605](#)

Gene Name MAP2K2

Gene Alias FLJ26075, MAPKK2, MEK2, MKK2, PRKMK2

Gene Description mitogen-activated protein kinase kinase 2

Omim ID [115150 601263](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene. [provided by RefSeq]

Other Designations

ERK activator kinase 2|MAP kinase kinase 2|MAPK/ERK kinase 2|dual specificity mitogen-activated protein kinase kinase 2|mitogen-activated protein kinase kinase 2, p45

Pathway

- [Acute myeloid leukemia](#)
- [B cell receptor signaling pathway](#)
- [Bladder cancer](#)
- [Chronic myeloid leukemia](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Gap junction](#)
- [Glioma](#)
- [GnRH signaling pathway](#)

- [Insulin signaling pathway](#)
- [Long-term depression](#)
- [Long-term potentiation](#)
- [MAPK signaling pathway](#)
- [Melanogenesis](#)
- [Melanoma](#)
- [Natural killer cell mediated cytotoxicity](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Pathways in cancer](#)
- [Prion diseases](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Renal cell carcinoma](#)
- [T cell receptor signaling pathway](#)
- [Thyroid cancer](#)
- [Toll-like receptor signaling pathway](#)
- [Vascular smooth muscle contraction](#)
- [VEGF signaling pathway](#)

Disease

- [Abnormalities](#)
- [Ectodermal Dysplasia](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)

- [Heart Defects](#)
- [LEOPARD Syndrome](#)
- [Mental Retardation](#)
- [Noonan Syndrome](#)
- [Skin Abnormalities](#)
- [Syndrome](#)