


Full-Length

PSAP (Human) Recombinant Protein (P02)

Catalog # H00005660-P02

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human PSAP full-length ORF (AAH04275, 1 a.a. - 524 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MYALFLLASLLGAALAGPVLGLKECTRGSAVWCQNVKTASDCGAVKHCLQTVWNKPTVKSLPC
DICKDVVTAAGDMLKDNATEEEILVYLEKTCDWLPKPNMSASCKENDSYLPVILDIKGEMSRPGE
VCSALNLCESLQKHLAELNHQKQLESNKIPELDITEVVAPFMANIPLLLYPQDGPRSKPQPKDNGD
VCQDCIQMVTDIQTAVRTNSTFVQALVEHVKEECDRLGPGMADICKNYISQYSEIAIQMMMHHMQPK
EICALVGFCDEVKEMPMQTLVPAKVASKNVIPALELVEPIKKHEVPAKSDVYCEVCEFLVKEVTK
LIDNNKTEKEILDAFDKMCSKLPKSLSEECQEVVDTYGSSILSILLEEVSPELVCSMLHLCSGTRLPL
ALTVHVTQPKDGGFCEVCKKLVGYLDRNLEKNSTKQEILAALEKGCFLPDYQKQCDQFVAEY
EPVLIEILVEVMDPSFVCLKIGACPLAHKPLLGTEKCMWGPSYWCQNTETAACNAVEHCKRHVW
N

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

83.16

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — PSAP

Entrez GeneID[5660](#)**GeneBank Accession#**[BC004275](#)**Protein Accession#**[AAH04275](#)**Gene Name**

PSAP

Gene Alias

FLJ00245, GLBA, MGC110993, SAP1

Gene Description

prosaposin

Omim ID[176801](#) [249900](#) [610539](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a highly conserved glycoprotein which is a precursor for 4 cleavage products: saposins A, B, C, and D. Each domain of the precursor protein is approximately 80 amino acid residues long with nearly identical placement of cysteine residues and glycosylation sites. Saposins A-D localize primarily to the lysosomal compartment where they facilitate the catabolism of glycosphingolipids with short oligosaccharide groups. The precursor protein exists both as a secretory protein and as an integral membrane protein and has neurotrophic activities. Mutations in this gene have been associated with Gaucher disease, Tay-Sachs disease, and metachromatic leukodystrophy. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq]

Other Designations

OTTHUMP00000019776|sphingolipid activator protein-1

Pathway

- [Lysosome](#)

Disease

- [Alzheimer Disease](#)
- [Genetic Predisposition to Disease](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)
- [Schizophrenia](#)
- [Tobacco Use Disorder](#)