

DNAxPAb

Hard-to-Find
Antibody

PSMD5 DNAxPab

Catalog # H00005711-W01P

Size 200 ug

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human PSMD5 DNA using DNAx™ Immune technology.
Technology	DNAx™ Immune
Immunogen	Full-length human DNA
Sequence	MAAQALALLREVARLEAPLEELRALHSVLQAVPLNELRQQAAELRLGPLFSLLNENHREKTTLCV SILERLLQAMEPVHVARNLRVDLQRGLIHPDDSVKILTLISQIGRIVENS DAVTEILNNAELLKQVYCIG GENLSVAKAAIKSLSRISLTQAGLEALFESNLLDDLKSVMTNDIVRYRVYELIIEISSVSPESLNYCT TSGLV TQLLRELTGEDVLVRATCIEMVTSLAYTHHGRQYLAQEGVIDQISNIIVGADSDPFSSFYLP GVKFFGNLAVMDSPQQICERYPIFVEKVFEMIESQDPTMIGVAVDTVGILGSNVEGKQVLQKTGTR FERLLMRIGHQSKNAPVELKIRCLDAISSLLYPPEQQTDDLRLMTESWFSLSRDPLELFRGISSQ PFPELHCAALKVFTAIANQPWAQKLMFNSPGFVEYVVD RSV EHDKASKDAKYELVKALANSKTIA EIFGNPNYLRRLTYLSEGPYYVKPVSTTAVEGAE
Host	Rabbit
Reactivity	Human
Purification	Protein A
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)
- Flow Cytometry (Transfected cell)

Gene Info — PSMD5

Entrez GeneID [5711](#)

GeneBank Accession# [NM_005047.2](#)

Protein Accession# [NP_005038.1](#)

Gene Name PSMD5

Gene Alias KIAA0072, MGC23145, S5B

Gene Description proteasome (prosome, macropain) 26S subunit, non-ATPase, 5

Omim ID [604452](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator base. [provided by RefSeq]

Other Designations

26S protease subunit S5 basic|26S proteasome non-ATPase regulatory subunit 5|26S proteasome subunit S5B|OTTHUMP00000021990|proteasome 26S non-ATPase subunit 5