

PTPN7 rabbit monoclonal antibody

Catalog # H00005778-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human PTPN7 peptide using ARM Technology.
Immunogen	A synthetic peptide of human PTPN7 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human PTPN7 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — PTPN7

Entrez GeneID [5778](#)

GeneBank Accession# [PTPN7](#)

Gene Name PTPN7

Gene Alias BPTP-4, HEPTP, LC-PTP, LPTP, PTPNI

Gene Description protein tyrosine phosphatase, non-receptor type 7

Omim ID [176889](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The noncatalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Two alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

OTTHUMP00000034115|dual specificity phosphatase 1|hematopoietic protein-tyrosine phosphatase|protein-tyrosine phosphatase, nonreceptor-type, stress induced

Pathway

- [MAPK signaling pathway](#)