


RAD51 (Human) IP-WB Antibody Pair

Catalog # H00005888-PW1

Size 1 Set

Applications



Immunoprecipitation of RAD51 transfected lysate using mouse monoclonal anti-RAD51 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with rabbit polyclonal anti-RAD51.

Specification

Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of RAD51 transfected lysate using mouse monoclonal anti-RAD51 and Protein A Magnetic Bead (U0007), and immunoblotted with rabbit polyclonal anti-RAD51.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: mouse monoclonal anti-RAD51 (300 ug) 2. Antibody pair for WB: rabbit polyclonal anti-RAD51 (50 ul)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- Immunoprecipitation-Western Blot

[Protocol Download](#)

Gene Info — RAD51

Entrez GeneID [5888](#)

Gene Name RAD51

Gene Alias BRCC5, HRAD51, HsRad51, HsT16930, RAD51A, RECA

Gene Description RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae)

Omim ID [114480 179617](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the RAD51 protein family. RAD51 family members are highly similar to bacterial RecA and Saccharomyces cerevisiae Rad51, and are known to be involved in the homologous recombination and repair of DNA. This protein can interact with the ssDNA-binding protein RPA and RAD52, and it is thought to play roles in homologous pairing and strand transfer of DNA. This protein is also found to interact with BRCA1 and BRCA2, which may be important for the cellular response to DNA damage. BRCA2 is shown to regulate both the intracellular localization and DNA-binding ability of this protein. Loss of these controls following BRCA2 inactivation may be a key event leading to genomic instability and tumorigenesis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

BRCA1/BRCA2-containing complex, subunit 5|DNA repair protein RAD51 homolog 1|RAD51 homolog protein|RAD51, S.cerevisiae, homolog of, A|RecA, E. coli, homolog of|RecA-like protein|recombination protein A

Pathway

- [Homologous recombination](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)

Disease

- [Acute Disease](#)
- [Adenocarcinoma](#)
- [Ataxia telangiectasia](#)
- [Brain Neoplasms](#)

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Chromosome Aberrations](#)
- [Colorectal Neoplasms](#)
- [DNA Damage](#)
- [DNA Repair-Deficiency Disorders](#)
- [Esophageal Neoplasms](#)
- [Fanconi Anemia](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Myelodysplastic Syndromes](#)
- [Neoplasms](#)
- [Neuroma](#)
- [Occupational Diseases](#)

- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic Neoplasms](#)
- [Precancerous Conditions](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Radiodermatitis](#)
- [Recurrence](#)
- [Stomach Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)