

RAN rabbit monoclonal antibody

Catalog # H00005901-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human RAN peptide using ARM Technology.
Immunogen	A synthetic peptide of human RAN is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human RAN peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — RAN

Entrez GeneID [5901](#)

GeneBank Accession# [RAN](#)

Gene Name RAN

Gene Alias ARA24, Gsp1, TC4

Gene Description RAN, member RAS oncogene family

Omim ID [601179](#)

Gene Ontology [Hyperlink](#)

Gene Summary

RAN (ras-related nuclear protein) is a small GTP binding protein belonging to the RAS superfamily that is essential for the translocation of RNA and proteins through the nuclear pore complex. The RAN protein is also involved in control of DNA synthesis and cell cycle progression. Nuclear localization of RAN requires the presence of regulator of chromosome condensation 1 (RCC1). Mutations in RAN disrupt DNA synthesis. Because of its many functions, it is likely that RAN interacts with several other proteins. RAN regulates formation and organization of the microtubule network independently of its role in the nucleus-cytosol exchange of macromolecules. RAN could be a key signaling molecule regulating microtubule polymerization during mitosis. RCC1 generates a high local concentration of RAN-GTP around chromatin which, in turn, induces the local nucleation of microtubules. RAN is an androgen receptor (AR) coactivator that binds differentially with different lengths of polyglutamine within the androgen receptor. Polyglutamine repeat expansion in the AR is linked to Kennedy's disease (X-linked spinal and bulbar muscular atrophy). RAN coactivation of the AR diminishes with polyglutamine expansion within the AR, and this weak coactivation may lead to partial androgen insensitivity during the development of Kennedy's disease. [provided by RefSeq]

Other Designations OK/SW-cl.81|RanGTPase|guanosine triphosphatase Ran|member RAS oncogene family|ras-related nuclear protein

Disease

- [Adenocarcinoma](#)
- [Carcinoma](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)

- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Kidney Neoplasms](#)
- [Lung Neoplasms](#)
- [Mouth Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Precancerous Conditions](#)
- [Premature Birth](#)