


RARA monoclonal antibody (M09A), clone 2C3

Catalog # H00005914-M09A


Size 200 uL

Applications



Western Blot (Cell lysate)

RARA monoclonal antibody (M09A), clone 2C3. Western Blot analysis of RARA expression in HeLa S3 NE (Cat # L013V3).



Western Blot detection against Immunogen (37.84 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant RARA.

Immunogen

RARA (NP_000955, 315 a.a. ~ 424 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence

QLLPLEMDDAETGLLSAICLICGDRQDLEQPDRVDMLQEPLLEALKVYVRKRRPSRPHMFPMKML
MKITDLRSISAKGAERVITLKMEIPGSMPPPLIQEMLLENSEGLDTLS

Host

Mouse

Reactivity

Human

Interspecies Antigen Sequence	Mouse (99); Rat (99)
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .
Storage Buffer	In ascites fluid
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

RARA monoclonal antibody (M09A), clone 2C3. Western Blot analysis of RARA expression in Hela S3 NE (Cat # L013V3).

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

Gene Info — RARA

Entrez GeneID	5914
GeneBank Accession#	NM_000964
Protein Accession#	NP_000955
Gene Name	RARA
Gene Alias	NR1B1, RAR
Gene Description	retinoic acid receptor, alpha
Omim ID	180240
Gene Ontology	Hyperlink

Gene Summary

Retinoid signaling is transduced by 2 families of nuclear receptors, retinoic acid receptor (RAR) and retinoid X receptor (RXR; see MIM 180245), which form RXR/RAR heterodimers. In the absence of ligand, DNA-bound RXR/RARA represses transcription by recruiting the corepressors NCO R1 (MIM 600849), SMRT (NCOR2; MIM 600848), and histone deacetylase (see MIM 601241). When ligand binds to the complex, it induces a conformational change allowing the recruitment of coactivators, histone acetyltransferases (see MIM 603053), and the basic transcription machinery. Translocations that always involve rearrangement of the RARA gene are a cardinal feature of acute promyelocytic leukemia (APL; MIM 612376). The most frequent translocation is t(15,17)(q21;q22), which fuses the RARA gene with the PML gene (MIM 102578) (Vitoux et al., 2007 [PubMed 174 68032]).[supplied by OMIM]

Other Designations

OTTHUMP00000164454|OTTHUMP00000164456|Retinoic acid receptor, alpha polypeptide|nucleophosmin-retinoic acid receptor alpha fusion protein NPM-RAR long form

Pathway

- [Acute myeloid leukemia](#)
- [Pathways in cancer](#)

Disease

- [Alcoholism](#)
- [Alzheimer disease](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Bipolar Disorder](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Drug Toxicity](#)
- [Edema](#)

- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Liver Cirrhosis](#)
- [Mental Disorders](#)
- [Metabolic Syndrome X](#)
- [Myopia](#)
- [Neoplasms](#)
- [Osteoporosis](#)
- [Schizophrenia](#)