

# RPL13 rabbit monoclonal antibody

Catalog # H00006137-K

Size 100 ug x up to 3

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human RPL13 peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human RPL13 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human RPL13 peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — RPL13

Entrez GeneID	<a href="#">6137</a>
GeneBank Accession#	<a href="#">RPL13</a>
Gene Name	RPL13
Gene Alias	BBC1, D16S444E, FLJ27453, FLJ27454, MGC117342, MGC71373
Gene Description	ribosomal protein L13
Omim ID	<a href="#">113703</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L13E family of ribosomal proteins. It is located in the cytoplasm. This gene is expressed at significantly higher levels in benign breast lesions than in breast carcinomas. Transcript variants derived from alternative splicing and/or alternative polyadenylation exist; these variants encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq]
Other Designations	60S ribosomal protein L13 OK/SW-cl.46 OTTHUMP00000165256 breast basic conserved protein 1

## Pathway

- [Ribosome](#)