


Full-Length

S100A9 (Human) Recombinant Protein (P01)

Catalog # H00006280-P01 Size 25 ug, 10 ug

Applications



Specification	
Product Description	Human S100A9 full-length ORF (AAH47681, 1 a.a 114 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MTCKMSQLERNIETIINTFHQYSVKLGHPDTLNQGEFKELVRKDLQNFLKKENRNEKVIEHIMEDLD TNADKQLSFEEFIMLMARLTWASHEKMHEGDEGPGHHHKPGLGEGTP
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	38.28
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.



Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — S100A9	
Entrez GenelD	6280
GeneBank Accession#	BC047681
Protein Accession#	AAH47681
Gene Name	S100A9
Gene Alias	60B8AG, CAGB, CFAG, CGLB, L1AG, LIAG, MAC387, MIF, MRP14, NIF, P14
Gene Description	S100 calcium binding protein A9
Omim ID	123886
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-han d calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide ra nge of cells, and involved in the regulation of a number of cellular processes such as cell cycle pro gression and differentiation. S100 genes include at least 13 members which are located as a clus ter on chromosome 1q21. This protein may function in the inhibition of casein kinase and altered expression of this protein is associated with the disease cystic fibrosis. [provided by RefSeq
Other Designations	OTTHUMP00000015331 S100 calcium-binding protein A9 S100 calcium-binding protein A9 (cal granulin B) calgranulin B

Publication Reference



Methods for predicting rheumatoid arthritis treatment response.

Olivier Vittecoq, Thierry Lequerre, Pascal Cosette, Olivier Boyer, Xavier Le Loet, Julie Hardouin, Antoine Obry.

United States Patent Application Publication 2016 May; [Epub].

Application: ELISA, Human, Serum

Melanocyte and melanoma cell activation by calprotectin.

Shirley SH, von Maltzan K, Robbins PO, Kusewitt DF.

Journal of Skin Cancer 2014 Aug; 2014:846249.

Application: Cell Proliferation Assay, Human, Melanocyte, Melanoma cells

 CD14(+)S100A9(+) monocytic myeloid-derived suppressor cells and their clinical relevance in non-small cell lung cancer.

Feng PH, Lee KY, Chang YL, Chan YF, Kuo LW, Lin TY, Chung FT, Kuo CS, Yu CT, Lin SM, Wang CH, Chou CL, Huang CD, Kuo HP.

American Journal of Respiratory and Critical Care Medicine 2012 Nov; 186(10):1025.

Application: Func, Human, A-549 cells

In vivo targeting of inflammation-associated myeloid-related protein 8/14 via gadolinium immunonanoparticles.

Maiseyeu A, Badgeley MA, Kampfrath T, Mihai G, Deiuliis JA, Liu C, Sun Q, Parthasarathy S, Simon DI, Croce K, Rajagopalan S.

Arteriosclerosis, Thrombosis, and Vascular Biology 2012 Apr; 32(4):962.

 Lack of an Endogenous Anti-inflammatory Protein in Mice Enhances Colonization of B16F10 Melanoma Cells in the Lungs.

Saha A, Lee YC, Zhang Z, Chandra G, Su SB, Mukherjee AB.

The Journal of Biological Chemistry 2010 Apr; 285(14):10822.

Application: Func, Mouse, B16F10 cells

Cationic polypeptides are required for anti-HIV-1 activity of human vaginal fluid.

Venkataraman N, Cole AL, Svoboda P, Pohl J, Cole AM.

Journal of Immunology 2005 Dec; 175(11):7560.

Application: WB, Human, Human vaginal fluid

Disease

- Dermatitis
- Genetic Predisposition to Disease



• Kidney Calculi