

CCL17 rabbit monoclonal antibody

Catalog # H00006361-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human CCL17 peptide using ARM Technology.
Immunogen	A synthetic peptide of human CCL17 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human CCL17 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — CCL17

Entrez GeneID	6361
GeneBank Accession#	CCL17
Gene Name	CCL17
Gene Alias	A-152E5.3, ABCD-2, MGC138271, MGC138273, SCYA17, TARC
Gene Description	chemokine (C-C motif) ligand 17
Omim ID	601520
Gene Ontology	Hyperlink
Gene Summary	This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells. [provided by RefSeq]
Other Designations	OTTHUMP00000164673 T cell-directed CC chemokine small inducible cytokine A17 small inducible cytokine subfamily A (Cys-Cys), member 17 thymus and activation-regulated chemokine

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)

Disease

- [Asthma](#)
- [Bronchiolitis](#)
- [Dermatitis](#)
- [Eosinophilia](#)

- [Genetic Predisposition to Disease](#)
- [Infant](#)
- [Multiple Sclerosis](#)
- [Respiratory Syncytial Virus Infections](#)