

ST6GAL1 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00006480-T02 Size 100 uL

Applications



20

10

SDS-PAGE Gel

ST6GAL1 transfected lysate.

Western Blot

Lane 1: ST6GAL1 transfected lysate (20.80 KDa) Lane 2: Non-transfected lysate.

Specification	
Transfected Cell Line	293T
Plasmid	pCMV-ST6GAL1 full-length
Host	Human
Theoretical MW (kDa)	20.8
Quality Control Testing	Transient overexpression cell lysate was tested with Anti-ST6GAL1 antibody (<u>H00006480-D01P</u>) by Western Blots. SDS-PAGE Gel ST6GAL1 transfected lysate. Western Blot Lane 1: ST6GAL1 transfected lysate (20.80 KDa) Lane 2: Non-transfected lysate.



Product Information

Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

• Western Blot

Gene Info — ST6GAL1

<u>6480</u>
<u>NM_173217.1</u>
<u>NP_775324.1</u>
ST6GAL1
CD75, MGC48859, SIAT1, ST6Gall, ST6N
ST6 beta-galactosamide alpha-2,6-sialyltranferase 1
<u>109675</u>
Hyperlink
This gene encodes a member of glycosyltransferase family 29. The encoded protein is a type II m embrane protein that catalyzes the transfer of sialic acid from CMP-sialic acid to galactose-contai ning substrates. The protein, which is normally found in the Golgi but can be proteolytically proces sed to a soluble form, is involved in the generation of the cell-surface carbohydrate determinants a nd differentiation antigens HB-6, CD75, and CD76. This gene has been incorrectly referred to as CD75. Three transcript variants encoding two different isoforms have been described. [provided by RefSeq
CMP-N-acetylneuraminate beta-galactosamide alpha-2,6-sialyltransferase ST6Gal l alpha 2,6-ST sialyltransferase 1 (beta-galactoside alpha-2,6-sialyltransferase) sialyltransferase 1 (beta-galacto

Pathway

- Metabolic pathways
- <u>N-Glycan biosynthesis</u>



Disease

• Tobacco Use Disorder