


STAT3 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00006774-T01


Size 100 uL

Applications



SDS-PAGE Gel

STAT3 transfected lysate.



Western Blot

Lane 1: STAT3 transfected lysate (88.1 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-STAT3 full-length
Host	Human
Theoretical MW (kDa)	88.1
Interspecies Antigen Sequence	Mouse (99); Rat (99)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-STAT3 antibody ([H00006774-B01P](#)) by Western Blots.
SDS-PAGE Gel
STAT3 transfected lysate.
Western Blot
Lane 1: STAT3 transfected lysate (88.1 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — STAT3

Entrez GeneID

[6774](#)

GeneBank Accession#

[NM_139276](#)

Protein Accession#

[AAH14482.1](#)

Gene Name

STAT3

Gene Alias

APRF, FLJ20882, HIES, MGC16063

Gene Description

signal transducer and activator of transcription 3 (acute-phase response factor)

Omim ID

[102582](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations

DNA-binding protein APRF|acute-phase response factor|signal transducer and activator of transcription 3

Pathway

- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)

Disease

- [Abortion](#)
- [Asthma](#)
- [Autoimmune Diseases](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Depressive Disorder](#)

- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [DNA Damage](#)
- [Eczema](#)
- [Edema](#)
- [Fatty Liver](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hepatitis C](#)
- [Infant](#)
- [Inflammation](#)
- [Inflammatory Bowel Diseases](#)
- [Insulin Resistance](#)
- [Kidney Neoplasms](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neoplasm Metastasis](#)

- [Obesity](#)
- [Ovarian Neoplasms](#)
- [Pulmonary Disease](#)
- [Rectal Fistula](#)
- [Respiratory Syncytial Virus Infections](#)
- [Spondylitis](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tooth Abnormalities](#)