


Full-Length

# SURF4 (Human) Recombinant Protein (P01)

Catalog # H00006836-P01 Size 25 ug, 10 ug

## **Applications**



Specification	
Product Description	Human SURF4 full-length ORF ( NP_149351.1, 1 a.a 269 a.a.) recombinant protein with GST-tag a t N-terminal.
Sequence	MGQNDLMGTAEDFADQFLRVTKQYLPHVARLCLISTFLEDGIRMWFQWSEQRDYIDTTWNCGYLL ASSFVFLNLLGQLTGCVLVLSRNFVQYACFGLFGIIALQTIAYSILWDLKFLMRNLALGGGLLLLLAE SRSEGKSMFAGVPTMRESSPKQYMQLGGRVLLVLMFMTLLHFDASFFSIVQNIVGTALMILVAIGF KTKLAALTLVVWLFAINVYFNAFWTIPVYKPMHDFLKYDFFQTMSVIGGLLLVVALGPGGVSMDEK KKEW
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	56.8
Interspecies Antigen Sequence	Mouse (99); Rat (99)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCI, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.



#### **Product Information**

Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

### **Applications**

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — SURF4	
Entrez GenelD	<u>6836</u>
GeneBank Accession#	NM_033161.2
Protein Accession#	NP_149351.1
Gene Name	SURF4
Gene Alias	ERV29, FLJ22993, MGC102753
Gene Description	surfeit 4
Omim ID	<u>185660</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene is located in the surfeit gene cluster, which is comprised of very tightly linked housekee ping genes that do not share sequence similarity. The encoded protein is a conserved integral me mbrane protein containing multiple putative transmembrane regions. In eukaryotic cells, protein transport between the endoplasmic reticulum and Golgi compartments is mediated in part by non-clathrin-coated vesicular coat proteins (COPs). The specific function of this protein has not been determined but its yeast homolog is directly required for packaging glycosylated pro-alpha-factor in to COPII vesicles. This gene uses multiple polyadenylation sites, resulting in transcript length variation. The existence of alternatively spliced transcript variants has been suggested, but their validity has not been determined. [provided by RefSeq
Other Designations	OTTHUMP00000022476 surface 4 integral membrane protein surfeit locus protein 4



### Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Tobacco Use Disorder