


MaxPab®

TAF10 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00006881-D01P

Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of TAF10 expression in transfected 293T cell line ([H00006881-T01](#)) by TAF10 MaxPab polyclonal antibody.

Lane 1: TAF10 transfected lysate(21.70 kDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human TAF10 protein.
Immunogen	TAF10 (NP_006275.1, 1 a.a. ~ 218 a.a) full-length human protein.
Sequence	MSCSGSGADPEAAPASAASAPGPAPPVSAPAALPSSTAAENKASPAGTAGGPGAGAAAGGTG PLAARAGEPAERRGAAPVSAGGAAPPEGAIISNGVYVLPSAANGDVKPVVSSSTPLVDFLMQLEDY TPTIPDAVTGYLNRAGFEASDPRIIRLISLAAQKFISDIANDALQHCKMKGTASGSSRSKSKDRKYT LTMEDLTPALSEYGINVKKPHYFT
Host	Rabbit
Reactivity	Human
Interspecies Antigen Sequence	Mouse (92); Rat (93)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of TAF10 expression in transfected 293T cell line ([H00006881-T01](#)) by TAF10 MaxPab polyclonal antibody.

Lane 1: TAF10 transfected lysate(21.70 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — TAF10

Entrez GeneID [6881](#)

GeneBank Accession# [NM_006284.2](#)

Protein Accession# [NP_006275.1](#)

Gene Name TAF10

Gene Alias TAF2A, TAF2H, TAFII30

Gene Description TAF10 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 30kDa

Omim ID [600475](#)

Gene Ontology [Hyperlink](#)

Gene Summary

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptide s. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the small subunits of TFIID that is associated with a subset of TFIID complexes. Studies with human and mammalian cells have shown that this subunit is required for transcriptional activation by the estrogen receptor, for progression through the cell cycle, and may also be required for certain cellular differentiation programs . [provided by RefSeq]

Other Designations

TAF10 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 30 kDa|TATA box binding protein (TBP)-associated factor, RNA polymerase II, H, 30kDa|TBP-related factor 10|transcription initiation factor TFIID 30 kD subunit

Pathway

- [Basal transcription factors](#)