


TAF11 monoclonal antibody (M04), clone 2G9


Catalog # H00006882-M04 Size 100 ug

Applications




Western Blot (Cell lysate)

TAF11 monoclonal antibody (M04), clone 2G9 Western Blot analysis of TAF11 expression in Hela S3 NE (Cat # L013V3).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)


Immunoperoxidase of monoclonal antibody to TAF11 on formalin-fixed paraffinembedded human placenta. [antibody concentration 3 ug/ml]



Sandwich ELISA (Recombinant protein)


Detection limit for recombinant GST tagged TAF11 is approximately 0.1ng/ml as a capture antibody.





Immunofluorescence

Immunofluorescence of monoclonal antibody to TAF11 on HeLa cell. [antibody concentration 10 ug/ml]



Western Blot detection against Immunogen (31.46 KDa).

Specification	
Product Description	Mouse monoclonal antibody raised against a partial recombinant TAF11.
Immunogen	TAF11 (NP_005634, 158 a.a. \sim 210 a.a) partial recombinant protein with GST tag. MW of the GST t ag alone is 26 KDa.
Sequence	SKVFVGEVVEEALDVCEKWGEMPPLQPKHMREAVRRLKSKGQIPNSKHKKIIF
Host	Mouse
Reactivity	Human
Isotype	lgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (31.46 KDa).
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications



Western Blot (Cell lysate)

 $TAF11\ monoclonal\ antibody\ (M04),\ clone\ 2G9\ Western\ Blot\ analysis\ of\ TAF11\ expression\ in\ Hela\ S3\ NE\ (\ Cat\ \#\ L013V3\).$

Protocol Download

Western Blot (Recombinant protein)

Protocol Download

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunoperoxidase of monoclonal antibody to TAF11 on formalin-fixed paraffin-embedded human placenta. [antibody concentration 3 ug/ml]

Protocol Download

Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged TAF11 is approximately 0.1ng/ml as a capture antibody.

Protocol Download

- ELISA
- Immunofluorescence

Immunofluorescence of monoclonal antibody to TAF11 on HeLa cell. [antibody concentration 10 ug/ml]

Gene Info — TAF11	
Entrez GeneID	<u>6882</u>
GeneBank Accession#	NM_005643
Protein Accession#	NP_005634
Gene Name	TAF11
Gene Alias	MGC:15243, PRO2134, TAF2I, TAFII28
Gene Description	TAF11 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 28kDa
Omim ID	600772
Gene Ontology	<u>Hyperlink</u>



Product Information

Gene Summary

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptide s. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the rem ainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit of TFIID that is present in all TFIID complexes and interacts with TBP. This subunit also interacts with another small subunit, TAF13, to form a heterodimer with a structure similar to the histone core structure. [provided by RefSeq

Other Designations

OTTHUMP00000016241|TAF11 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 28 kD|TATA box binding protein (TBP)-associated factor, RNA polymerase II, I, 28kD|TBP-associated factor 11|TFIID subunit p30-beta|transcription initiation facto

Pathway

Basal transcription factors

Disease

Kidney Failure