

TAF13 rabbit monoclonal antibody

Catalog # H00006884-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human TAF13 peptide using ARM Technology.
Immunogen	A synthetic peptide of human TAF13 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human TAF13 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — TAF13

Entrez GeneID	6884
GeneBank Accession#	TAF13
Gene Name	TAF13
Gene Alias	MGC22425, TAF2K, TAFII18
Gene Description	TAF13 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 18kDa
Omim ID	600774
Gene Ontology	Hyperlink
Gene Summary	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptide s. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit associated with a subset of TFIID complexes. This subunit interacts with TBP and with two other small subunits of TFIID, TAF10 and TAF11. There is a pseudogene located on chromosome 6. [provided by RefSeq]
Other Designations	OTTHUMP00000039344 TATA box binding protein (TBP)-associated factor, RNA polymerase II, K, 18kD TBP-associated factor 13 transcription initiation factor TFIID 18 kD subunit

Pathway

- [Basal transcription factors](#)