

DNAxPAb

Hard-to-Find
Antibody

TBCC DNAxPab

Catalog # H00006903-W01P

Size 200 ug

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human TBCC DNA using DNAx™ Immune technology.
Technology	DNAx™ Immune
Immunogen	Full-length human DNA
Sequence	MESVSCSAAAVRTGDMESQRDLSLPERLQRREQERQLEVERRKQKRQNQEVEKENSHEFFVATFARERAAVEELLERAESVERLEEAASRLQGLQKLINDSVFFLAAYDLRQGQEALARLQAALAERRRGLQPKKRFAFKTRGKDAASSTKVDAAPGIPPAVESIQDSPLPKKAEGDLGPSWVCGFSNLESQVLEKRASELHQRDVLLTELSNCTVRLYGPNPTLRLTKAHSCKLLCGPVSTSVFLED CSDCVLAVACQQLRIHSTKDTRIFLQVTSRAVEDCSGIQFAPYTWSYPEIDKDFESSGLDRSKNNWNDVDDFNWLARDMASPNWSILPEEERNIQWD
Host	Rabbit
Reactivity	Human
Purification	Protein A
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)

- Flow Cytometry (Transfected cell)

Gene Info — TBCC

Entrez GeneID	6903
GeneBank Accession#	NM_003192.1
Protein Accession#	NP_003183.1
Gene Name	TBCC
Gene Alias	CFC
Gene Description	tubulin folding cofactor C
Omim ID	602971
Gene Ontology	Hyperlink
Gene Summary	Cofactor C is one of four proteins (cofactors A, D, E, and C) involved in the pathway leading to correctly folded beta-tubulin from folding intermediates. Cofactors A and D are believed to play a role in capturing and stabilizing beta-tubulin intermediates in a quasi-native confirmation. Cofactor E binds to the cofactor D/beta-tubulin complex; interaction with cofactor C then causes the release of beta-tubulin polypeptides that are committed to the native state. [provided by RefSeq]
Other Designations	OTTHUMP00000016406 beta-tubulin cofactor C tubulin-specific chaperone c