


TRAF6 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00007189-T02


Size 100 uL

Applications



SDS-PAGE Gel

TRAF6 transfected lysate.



Western Blot

Lane 1: TRAF6 transfected lysate (59.6 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-TRAF6 full-length

Host Human

Theoretical MW (kDa) 59.6

Quality Control Testing Transient overexpression cell lysate was tested with Anti-TRAF6 antibody ([H00007189-D01](#)) by Western Blots.
 SDS-PAGE Gel
 TRAF6 transfected lysate.
 Western Blot
 Lane 1: TRAF6 transfected lysate (59.6 KDa)
 Lane 2: Non-transfected lysate.

Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — TRAF6

Entrez GeneID	7189
GeneBank Accession#	BC031052
Protein Accession#	AAH31052.1
Gene Name	TRAF6
Gene Alias	MGC:3310, RNF85
Gene Description	TNF receptor-associated factor 6
Omim ID	602355
Gene Ontology	Hyperlink

Gene Summary	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins are associated with, and mediate signal transduction from members of the TNF receptor superfamily. This protein mediates the signaling not only from the members of the TNF receptor superfamily, but also from the members of the Toll/IL-1 family. Signals from receptors such as CD40, TNFSF11/RANCE and IL-1 have been shown to be mediated by this protein. This protein also interacts with various protein kinases including IRAK1/IRAK, SRC and PKCzeta, which provides a link between distinct signaling pathways. This protein functions as a signal transducer in the NF-kappaB pathway that activates I kappaB kinase (IKK) in response to proinflammatory cytokines. The interaction of this protein with UBE2N/UBC13, and UBE2V1/UEV1A, which are ubiquitin conjugating enzymes catalyzing the formation of polyubiquitin chains, has been found to be required for IKK activation by this protein. Two alternatively spliced transcript variants encoding identical proteins have been reported. [provided by RefSeq]
---------------------	---

Other Designations	-
---------------------------	---

Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [Toll-like receptor signaling pathway](#)
- [Ubiquitin mediated proteolysis](#)

Disease

- [Alzheimer disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)

- [Hypersensitivity](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Narcolepsy](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Thyroid Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)