UBA7 rabbit monoclonal antibody

Catalog # H00007318-K

ocification

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human UBA7 peptide using ARM Technology.
Immunogen	A synthetic peptide of human UBA7 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human UBA7 peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download



• ELISA

Gene Info — UBA7	
Entrez GenelD	<u>7318</u>
GeneBank Accession#	UBA7
Gene Name	UBA7
Gene Alias	D8, MGC12713, UBA1B, UBE1L, UBE2
Gene Description	ubiquitin-like modifier activating enzyme 7
Omim ID	<u>191325</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnor mal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzym es: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-prot ein ligases, or E3s. This gene encodes a member of the E1 ubiquitin-activating enzyme family. Th e encoded enzyme is a retinoid target that triggers promyelocytic leukemia (PML)/retinoic acid re ceptor alpha (RARalpha) degradation and apoptosis in acute promyelocytic leukemia, where it is involved in the conjugation of the ubiquitin-like interferon-stimulated gene 15 protein. [provided by RefSeq
Other Designations	UBA1, ubiquitin-activating enzyme E1 homolog B UBA7, ubiquitin-activating enzyme E1 ubiquitin- activating enzyme E1-like ubiquitin-activating enzyme E1-related protein ubiquitin-activating enzy me-2

Pathway

<u>Ubiquitin mediated proteolysis</u>

Disease

- <u>Crohn Disease</u>
- Genetic Predisposition to Disease