

USF2 monoclonal antibody (M01), clone 5E9

Catalog # H00007392-M01

Size 100 ug

Applications




Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in HeLa.




Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in different cell lines.




Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in HeLa S3 NE (Cat # L013V3).



Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in PC-12 (Cat # L012V1).




Western Blot (Transfected lysate)

Western Blot analysis of USF2 expression in transfected 293T cell line by USF2 monoclonal antibody (M01), clone 5E9.


Lane 1: USF2 transfected lysate (37 KDa).

Lane 2: Non-transfected lysate.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunoperoxidase of monoclonal antibody to USF2 on formalin-fixed paraffin-embedded human cerebral cortex. [antibody concentration 1 ug/ml]




Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged USF2 is approximately 0.3 ng/ml as a capture antibody.



Immunofluorescence

Immunofluorescence of monoclonal antibody to USF2 on HeLa cell. [antibody concentration 10 ug/ml]



Western Blot detection against Immunogen (36.74 KDa).

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant USF2.
Immunogen	USF2 (NP_003358, 1 a.a. ~ 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	MDMLDPGLDPAASATAAAAASHDKGPEAEEGVELQEGGDGPGAEEQTAVAITSVQQAAGFDH NIQYQFRTE TNGGQV TYRVVQVTDGQLDGQGDTAGAVS
Host	Mouse
Reactivity	Human, Rat
Interspecies Antigen Sequence	Mouse (96); Rat (96)
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in HeLa.

[Protocol Download](#)

- Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in different cell lines.

[Protocol Download](#)

- Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in Hela S3 NE (Cat # L013V3).

[Protocol Download](#)

- Western Blot (Cell lysate)

USF2 monoclonal antibody (M01), clone 5E9. Western Blot analysis of USF2 expression in PC-12 (Cat # L012V1).

[Protocol Download](#)

- Western Blot (Transfected lysate)

Western Blot analysis of USF2 expression in transfected 293T cell line by USF2 monoclonal antibody (M01), clone 5E9.

Lane 1: USF2 transfected lysate(37 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunoperoxidase of monoclonal antibody to USF2 on formalin-fixed paraffin-embedded human cerebral cortex. [antibody concentration 1 ug/ml]

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged USF2 is approximately 0.3ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

- Immunofluorescence

Immunofluorescence of monoclonal antibody to USF2 on HeLa cell. [antibody concentration 10 ug/ml]

Gene Info — USF2

Entrez GeneID [7392](#)

GeneBank Accession# [NM_003367](#)

Protein Accession# [NP_003358](#)

Gene Name USF2

Gene Alias FIP, bHLHb12

Gene Description	upstream transcription factor 2, c-fos interacting
Omim ID	600390
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidin e-rich initiator (Inr) elements and E-box motifs. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq]
Other Designations	c-fos interacting protein upstream stimulatory factor 2

Disease

- [Alzheimer disease](#)
- [Biliary Atresia](#)
- [Cardiovascular Diseases](#)
- [Choledochal Cyst](#)
- [Diabetes Complications](#)
- [Disease Progression](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Osteoporosis](#)