


HMGA2 monoclonal antibody (M01), clone 2D10

Catalog # H00008091-M01


Size 100 ug

Applications




Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged HMGA2 is approximately 0.1ng/ml as a capture antibody.



Immunofluorescence

Immunofluorescence of monoclonal antibody to HMGA2 on HeLa cell .
[antibody concentration 10 ug/ml]



Western Blot detection against Immunogen (35.86 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant HMGA2.

Immunogen	HMGA2 (NP_003474, 1 a.a. ~ 92 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	MSARGEAGAGQPSTSAQGQPAAPAPQKRGRGRPRKQQQEPTGEPSPKRPRGRPKGSKNKSPS KAAQKKAETGEKRPRGRPRKWPQQVVQKKP
Host	Mouse
Reactivity	Human
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.86 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged HMGA2 is approximately 0.1ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

- Immunofluorescence

Immunofluorescence of monoclonal antibody to HMGA2 on HeLa cell . [antibody concentration 10 ug/ml]

Gene Info — HMGA2

Entrez GeneID	8091
GeneBank Accession#	NM_003483
Protein Accession#	NP_003474
Gene Name	HMGA2

Gene Alias	BABL, HMGI-C, HMGIC, LIPO, STQTL9
Gene Description	high mobility group AT-hook 2
Omim ID	151900 600698
Gene Ontology	Hyperlink
Gene Summary	<p>This gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhancosome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq]</p>
Other Designations	High-mobility group protein HMGI-C high-mobility group (nonhistone chromosomal) protein isoform I-C

Disease

- [Brain Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Insulin Resistance](#)
- [Leiomyoma](#)
- [Osteoarthritis](#)
- [Uterine Neoplasms](#)
- [Ventricular Dysfunction](#)