


HIST1H2AK (Human) Recombinant Protein (Q01)

Catalog # H00008330-Q01 Size 25 ug, 10 ug

Applications



| Specification | |
|----------------------------------|---|
| Product Description | Human HIST1H2AK partial ORF (NP_003501, 25 a.a 96 a.a.) recombinant protein with GST-tag a t N-terminal. |
| Sequence | QFPVGRVHRLLRKGNYAERVGAGAPVYLAAVLEYLTAEILELAGNAARDNKKTRIIPRHLQLAIRND EELNK |
| Host | Wheat Germ (in vitro) |
| Theoretical MW (kDa) | 33.66 |
| Interspecies Antigen Sequence | Mouse (99); Rat (96) |
| Preparation Method | in vitro wheat germ expression system |
| Purification | Glutathione Sepharose 4 Fast Flow |
| Quality Control Testing | 12.5% SDS-PAGE Stained with Coomassie Blue. |
| Storage Buffer | 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer. |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |
| Note | Best use within three months from the date of receipt of this protein. |



Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

| Gene Info — HIST1H2AK | |
|-----------------------|---|
| Entrez GenelD | 8330 |
| GeneBank Accession# | NM_003510 |
| Protein Accession# | NP_003501 |
| Gene Name | HIST1H2AK |
| Gene Alias | H2A/d, H2AFD |
| Gene Description | histone cluster 1, H2ak |
| Omim ID | 602788 |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, an d H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and f unctions in the compaction of chromatin into higher order structures. This gene is intronless and e ncodes a member of the histone H2A family. Transcripts from this gene lack polyA tails but instea d contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq |
| Other Designations | H2A histone family, member D OTTHUMP00000018006 histone 1, H2ak |

Pathway

Systemic lupus erythematosus