CDC14B rabbit monoclonal antibody

Catalog # H00008555-K

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human CDC14B peptide using ARM Technology.
Immunogen	A synthetic peptide of human CDC14B is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human CDC14B peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download

• ELISA

Gene Info — CDC14B

e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phe sphatase involved in the exit of cell mitosis and initiation of DNA replication, which suggests the r ole in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this ge ne results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq		
Gene Name CDC14B Gene Alias CDC14B3, Cdc14B1, Cdc14B2, hCDC14B Gene Description CDC14 cell division cycle 14 homolog B (S. cerevisiae) Omim ID 603505 Gene Ontology Hyperlink Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas e nole in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Entrez GenelD	<u>8555</u>
Gene Alias CDC14B3, Cdc14B1, Cdc14B2, hCDC14B Gene Description CDC14 cell division cycle 14 homolog B (S. cerevisiae) Omim ID 603505 Gene Ontology Hyperlink Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas e no cle in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	GeneBank Accession#	CDC14B
Gene Description CDC14 cell division cycle 14 homolog B (S. cerevisiae) Omim ID 603505 Gene Ontology Hyperlink Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas e no le in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Gene Name	CDC14B
Omim ID 603505 Gene Ontology Hyperlink Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatas involved in the exit of cell mitosis and initiation of DNA replication, which suggests the role in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Gene Alias	CDC14B3, Cdc14B1, Cdc14B2, hCDC14B
Gene Ontology Hyperlink Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatase e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatase involved in the exit of cell mitosis and initiation of DNA replication, which suggests the role in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Gene Description	CDC14 cell division cycle 14 homolog B (S. cerevisiae)
Gene Summary The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phosphatase involved in the exit of cell mitosis and initiation of DNA replication, which suggests the role in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this gene results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Omim ID	<u>603505</u>
e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine phe sphatase involved in the exit of cell mitosis and initiation of DNA replication, which suggests the r ole in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this ge ne results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq	Gene Ontology	Hyperlink
Other Designations CDC14 homolog B OTTHUMP00000021729 OTTHUMP00000021730 OTTHUMP00000063774	Gene Summary	The protein encoded by this gene is a member of the dual specificity protein tyrosine phosphatas e family. This protein is highly similar to Saccharomyces cerevisiae Cdc14, a protein tyrosine pho sphatase involved in the exit of cell mitosis and initiation of DNA replication, which suggests the r ole in cell cycle control. This protein has been shown to interact with and dephosphorylates tumor suppressor protein p53, and is thought to regulate the function of p53. Alternative splice of this ge ne results in 3 transcript variants encoding distinct isoforms. [provided by RefSeq
	Other Designations	CDC14 homolog B OTTHUMP00000021729 OTTHUMP00000021730 OTTHUMP00000063774

Pathway

• Cell cycle