


B4GALT4 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00008702-T01 Size 100 uL

Applications



SDS-PAGE Gel

B4GALT4 transfected lysate.

Western Blot

Lane 1: B4GALT4 transfected lysate (37.95 KDa) Lane 2: Non-transfected lysate.

| Specification | |
|----------------------------------|--------------------------|
| Transfected Cell Line | 293T |
| Plasmid | pCMV-B4GALT4 full-length |
| Host | Human |
| Theoretical MW (kDa) | 37.95 |
| Interspecies Antigen Sequence | Mouse (84); Rat (85) |



Product Information

| Quality Control Testing | Transient overexpression cell lysate was tested with Anti-B4GALT4 antibody (H00008702-B01) by | | |
|-------------------------|--|--|--|
| | Western Blots. | | |
| | SDS-PAGE Gel | | |
| | B4GALT4 transfected lysate. | | |
| | Western Blot | | |
| | Lane 1: B4GALT4 transfected lysate (37.95 KDa) | | |
| | Lane 2: Non-transfected lysate. | | |
| Storage Buffer | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue) | | |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. | | |
| | | | |

Applications

• Western Blot

Gene Info — B4GALT4

| Entrez GenelD | 8702 |
|---------------------|---|
| GeneBank Accession# | <u>BC004523.2</u> |
| Protein Accession# | <u>AAH04523.1</u> |
| Gene Name | B4GALT4 |
| Gene Alias | B4Gal-T4, beta4Gal-T4 |
| Gene Description | UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 4 |
| Omim ID | <u>604015</u> |
| Gene Ontology | Hyperlink |
| Gene Summary | This gene is one of seven beta-1,4-galactosyltransferase (beta4GaIT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate U DP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GaIT has a distinct function in the biosynthesis of different glycoconjugates an d saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signa I sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to fun ction as a transmembrane anchor. By sequence similarity, the beta4GaITs form four groups: beta 4GaIT1 and beta4GaIT2, beta4GaIT3 and beta4GaIT4, beta4GaIT5 and beta4GaIT6, and beta4GaIT7. The enzyme encoded by this gene appears to mainly play a role in glycolipid biosynthesis. T wo alternatively spliced transcript variants have been found for this gene. [provided by RefSeq |



Product Information

Other Designations

UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase 4|beta-N-acetylglucosaminyl-glycolipid beta -1,4-galactosyltransferase 4

Pathway

- Glycosphingolipid biosynthesis lacto and neolacto series
- Keratan sulfate biosynthesis
- Metabolic pathways