

EED rabbit monoclonal antibody

Catalog # H00008726-K Size 100 ug x up to 3

Rabbit monoclonal antibody raised against a human EED peptide using ARM Technology.
A synthetic peptide of human EED is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Rabbit
Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Overexpression vector and transfection into 293H cell line.
Human
Protein A
lgG
Antibody reactive against human EED peptide by ELISA and mammalian transfected lysate by West ern Blot.
In 1x PBS, pH 7.4
Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — EED	
Entrez GenelD	<u>8726</u>
GeneBank Accession#	<u>EED</u>
Gene Name	EED
Gene Alias	HEED, WAΠ1
Gene Description	embryonic ectoderm development
Omim ID	605984
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the Polycomb-group (PcG) family. PcG family members form multimeric protein complexes, which are involved in maintaining the transcriptional repressive state of genes over successive cell generations. This protein interacts with enhancer of zeste 2, the cyto plasmic tail of integrin beta7, immunodeficiency virus type 1 (HIV-1) MA protein, and histone deac etylase proteins. This protein mediates repression of gene activity through histone deacetylation, and may act as a specific regulator of integrin function. Two transcript variants encoding distinct is oforms have been identified for this gene. [provided by RefSeq
Other Designations	WD protein associating with integrin cytoplasmic tails 1