


# FADD (Human) Matched Antibody Pair

Catalog # H00008772-AP11

Size 1 Set

## Applications



Sandwich ELISA detection sensitivity ranging from 10 ng/ml to 100 ng/ml.

## Specification

<b>Product Description</b>	This antibody pair set comes with a matched antibody pair to detect and quantify the protein level of human FADD.
<b>Reactivity</b>	Human
<b>Interspecies Antigen Sequence</b>	Mouse (68); Rat (68)
<b>Quality Control Testing</b>	Standard curve using recombinant protein ( H00008772-P01 ) as an analyte. Sandwich ELISA detection sensitivity ranging from 10 ng/ml to 100 ng/ml.
<b>Supplied Product</b>	Antibody pair set content: 1. Capture antibody: rabbit MaxPab® affinity purified polyclonal anti-FADD (100 ug) 2. Detection antibody: mouse monoclonal anti-FADD, IgG1 Lambda (20 ug) *Reagents are sufficient for at least 1-2 x 96 well plates using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- ELISA Pair (Recombinant protein)

[Protocol Download](#)

## Gene Info — FADD

Entrez GeneID	<a href="#">8772</a>
Gene Name	FADD
Gene Alias	GIG3, MGC8528, MORT1
Gene Description	Fas (TNFRSF6)-associated via death domain
Omim ID	<a href="#">602457</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development. [provided by RefSeq]
Other Designations	Fas-associated via death domain Fas-associating death domain-containing protein Fas-associating protein with death domain growth-inhibiting gene 3 protein mediator of receptor-induced toxicity

## Pathway

- [Apoptosis](#)
- [Pathways in cancer](#)
- [Toll-like receptor signaling pathway](#)

## Disease

- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)

- [Hodgkin Disease](#)
- [Lupus Erythematosus](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)