


TNFRSF10B (Human) Recombinant Protein (Q01)

Catalog # H00008795-Q01

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human TNFRSF10B partial ORF (AAH01281, 71 a.a. - 170 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	QKRSSPSEGLCPPGHHISEDGRDCISCKYGQDYSTHWNDLLFCLRCTRCDSGEVELSPCTTTRN TVCQCEEFTFREEDSPEMCRKCRTGCPRGMVKVGDC
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	36.63
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — TNFRSF10B

Entrez GeneID	8795
GeneBank Accession#	BC001281
Protein Accession#	AAH01281
Gene Name	TNFRSF10B
Gene Alias	CD262, DR5, KILLER, KILLER/DR5, TRAIL-R2, TRAILR2, TRICK2, TRICK2A, TRICK2B, TRICK B, ZTNFR9
Gene Description	tumor necrosis factor receptor superfamily, member 10b
Omim ID	275355 603612
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. [provided by RefSeq]
Other Designations	Fas-like protein OTTHUMP00000123492 TNF-related apoptosis-inducing ligand receptor 2 TRAIL receptor 2 apoptosis inducing protein TRICK2A/2B apoptosis inducing receptor TRAIL-R2 cytotoxic TRAIL receptor-2 death domain containing receptor for TRAIL/Apo-2L d

Pathway

- [Apoptosis](#)
- [Cytokine-cytokine receptor interaction](#)

- [Natural killer cell mediated cytotoxicity](#)
- [p53 signaling pathway](#)

Disease

- [Asthma](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Prostatic Neoplasms](#)
- [Recurrence](#)
- [Waldenstrom Macroglobulinemia](#)

- [Werner syndrome](#)