

NR1I2 rabbit monoclonal antibody

Catalog # H00008856-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human NR1I2 peptide using ARM Technology.
Immunogen	A synthetic peptide of human NR1I2 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human NR1I2 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — NR1I2

Entrez GeneID [8856](#)

GeneBank Accession# [NR1I2](#)

Gene Name NR1I2

Gene Alias BXR, ONR1, PAR, PAR1, PAR2, PARq, PRR, PXR, SAR, SXR

Gene Description nuclear receptor subfamily 1, group I, member 2

Omim ID [603065](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized. [provided by RefSeq]

Other Designations orphan nuclear receptor PXR|steroid and xenobiotic receptor

Disease

- [Anemia](#)
- [Asthma](#)
- [Birth Weight](#)
- [Bone Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)

- [Cholangitis](#)
- [Cholestasis](#)
- [Colitis](#)
- [Colon cancer](#)
- [Colonic Neoplasms](#)
- [Coronary Artery Disease](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Drug-Induced Liver Injury](#)
- [Edema](#)
- [Epilepsy](#)
- [Fatty Liver](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hearing Loss](#)
- [HIV Infections](#)
- [Inflammatory Bowel Diseases](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Leukopenia](#)
- [Liver Cirrhosis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Myocardial Infarction](#)
- [Neoplasm Recurrence](#)

- [Neoplasms](#)
- [Neutropenia](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Recurrence](#)
- [Respiratory Tract Neoplasms](#)
- [Tobacco Use Disorder](#)