

DNAxPab

Hard-to-Find
Antibody

MED17 DNAxPab

Catalog # H00009440-W01P

Size 200 ug

Specification

Product Description Rabbit polyclonal antibody raised against a full-length human MED17 DNA using DNAx™ Immune technology.

Technology [DNAx™ Immune](#)

Immunogen Full-length human DNA

Sequence MSGVRAVRISIESACEKQVHEVGLDGTETYPPLSMSQNLARLAQRIDFSQGSSEEEEEAGTEG
DAQDWPGAGSSADQDDEEGVVKFQPSLWPWDSVRNNLRSALTEMCVLYDVLSMRDKKFMTL
DPVSQDALPPKQNPQTLQLISKKSLAGAAQILLKGAERLTKSVTENQENKLQRDFNSELLRLRQ
HWKLRKVGDKILGDLSYRSAGSLFPHHGTFEVIKNTDLDLKKIPEDYCPLDVQIPSDLEGSAYIKV
SIQKQAPDIGDLGTVNLFKRPLPKSKPGSPHWQTKLEAAQNVLLCKEIFAQLSREAVQIKSQVPHI
VVKNIISQPFPSLQLSISLCHSSNDKKSQKFATEKQCPEDHLYVLEHNLHLLIREFHKQTLSSIMM
PHPASAPFGHKRMRLSGPQAFDKNEINSLQSSEGLLEKIIKQAKHIFLRSAAATIDSLASRIEDPQI
QAHWSNINDVYESSVKVLITSQGYEQICKSIQLQLNIGVEQIRVVHRDGRVITLSYQEQELQDFLLSQ
MSQHQVHAVVQQLAKVMGWQVLSFSNHVGLGPIESIGNASAITVASPSGDYAISVRNGPESGSKIM
VQFPRNQCKDLPKSDVLQDNKWSHLRGPFEVQWNKMEGRNFVYKMELLMSALSPCLL

Host Rabbit

Reactivity Human

Purification Protein A

Quality Control Testing Antibody reactive against mammalian transfected lysate.

Storage Buffer In 1x PBS, pH 7.4

Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)

- Flow Cytometry (Transfected cell)

Gene Info — MED17

Entrez GeneID [9440](#)

GeneBank Accession# [BC021101.1](#)

Protein Accession# [AAH21101.1](#)

Gene Name MED17

Gene Alias CRSP6, CRSP77, DRIP80, FLJ10812, TRAP80

Gene Description mediator complex subunit 17

Omim ID [603810](#)

Gene Ontology [Hyperlink](#)

Gene Summary The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e. g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. [provided by RefSeq]

Other Designations cofactor required for Sp1 transcriptional activation, subunit 6 (77kD)|cofactor required for Sp1 transcriptional activation, subunit 6, 77kDa|thyroid hormone receptor-associated protein, 80-KD subunit|vitamin D receptor interacting protein 80-kD