

DNAxPAb

Hard-to-Find
Antibody

MED26 DNAxPab

Catalog # H00009441-W01P

Size 200 ug

Specification

Product Description Rabbit polyclonal antibody raised against a full-length human MED26 DNA using DNAx™ Immune technology.

Technology [DNAx™ Immune](#)

Immunogen Full-length human DNA

Sequence MTAAPASPQQIRDRLQLAIDPQSNIRNMVAVLEVISSLEKYPITKEALEETRLGKLINDVRKKTKNEELAKRAKKLLRSWQKLIIEPAHQHEAALRGLAGATGSANGGAHNCRPEVGAAGPPRSIHDLKSRNDLQRLPGQRLDRLGSRKRRGDQRDLGHPGPPPKVSKASHDPLVPNSSPLPTNGISGSPESFASSLDGSGHAGPEGSRLERDENDKHSGKIPVNAVVRPHTSSPGLGKPPGKCLQPKASVLQQLDRVDETPGPPHPKGPPRCFSFPRNSRHEGSFARQQSLYAPKGSVPSPSPRPQALDATQVPSPLPLAQPS TPPVRRELLPSAESPVCWLEQPESHQRLAGPGCKAGLSPAEPILLSRAGFSPDSSKADSDAASSGGSDSKKKKRYRPRDYTVNLDGQVAEAGVKPVRLKERKLTDFPMTRQIKPLTQKEPVRADSPVHMEQQSRTELDKQEAKASLQSPFEQTNWKELSRNEIIQSYLSRQSSLLSSSGAQTPGAHHFMSEYLKQEESTRQGARQLHVLVPQSPPTDLPGLTREVTQDDLDRIQASQWPGVNGCQDTQGNWYDW TQCISLDPHGDDGRLNILPYVCLD

Host Rabbit

Reactivity Human

Purification Protein A

Quality Control Testing Antibody reactive against mammalian transfected lysate.

Storage Buffer In 1x PBS, pH 7.4

Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)

- Flow Cytometry (Transfected cell)

Gene Info — MED26

Entrez GeneID [9441](#)

GeneBank Accession# [NM_004831.3](#)

Protein Accession# [NP_004822.2](#)

Gene Name MED26

Gene Alias CRSP7, CRSP70

Gene Description mediator complex subunit 26

Omim ID [605043](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e. g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. [provided by RefSeq]

Other Designations

cofactor required for Sp1 transcriptional activation, subunit 7 (70kD)|cofactor required for Sp1 transcriptional activation, subunit 7, 70kDa