


MAP4K4 monoclonal antibody (M06), clone 3B9


Catalog # H00009448-M06 Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged MAP4K4 is approximately 0.1ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.63 KDa).

| Specification | |
|---------------------|--|
| Product Description | Mouse monoclonal antibody raised against a partial recombinant MAP4K4. |
| Immunogen | MAP4K4 (NP_663719, 611 a.a. ~ 710 a.a) partial recombinant protein with GST tag. MW of the GS T tag alone is 26 KDa. |
| Sequence | VHPALQRPAEPQVQWSHLASLKNNVSPVSRSHSFSDPSPKFAHHHLRSQDPCPPSRSEVLSQ SSDSKSEAPDPTQKAWSRSDSDEVPPRVPVRTTSRSPV |
| Host | Mouse |
| Reactivity | Human |



Product Information

| Interspecies Antigen Sequence | Mouse (95); Rat (89) |
|----------------------------------|--|
| Isotype | lgG1 Kappa |
| Quality Control Testing | Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) . |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |

Applications

- Western Blot (Recombinant protein)
 <u>Protocol Download</u>
- Sandwich ELISA (Recombinant protein)
 Detection limit for recombinant GST tagged MAP4K4 is approximately 0.1ng/ml as a capture antibody.
 <u>Protocol Download</u>
- ELISA

Gene Info — MAP4K4

| Entrez GenelD | <u>9448</u> |
|---------------------|--|
| GeneBank Accession# | <u>NM_145686</u> |
| Protein Accession# | <u>NP_663719</u> |
| Gene Name | MAP4K4 |
| Gene Alias | FLH21957, FLJ10410, FLJ20373, FLJ90111, HGK, KIAA0687, NIK |
| Gene Description | mitogen-activated protein kinase kinase kinase kinase 4 |
| Omim ID | <u>604666</u> |
| Gene Ontology | Hyperlink |



Product Information

Gene Summary

The protein encoded by this gene is a member of the serine/threonine protein kinase family. This kinase has been shown to specifically activate MAPK8/JNK. The activation of MAPK8 by this kin ase is found to be inhibited by the dominant-negative mutants of MAP3K7/TAK1, MAP2K4/MKK4, and MAP2K7/MKK7, which suggests that this kinase may function through the MAP3K7-MAP2K 4-MAP2K7 kinase cascade, and mediate the TNF-alpha signaling pathway. Alternatively spliced t ranscript variants encoding different isoforms have been identified. [provided by RefSeq

Other Designations

HPK/GCK-like kinase|hepatocyte progenitor kinase-like/germinal center kinase-like kinase

Pathway

• MAPK signaling pathway