

BAG3 rabbit monoclonal antibody

Catalog # H00009531-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human BAG3 peptide using ARM Technology.
Immunogen	A synthetic peptide of human BAG3 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human BAG3 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — BAG3

Entrez GeneID	9531
GeneBank Accession#	BAG3
Gene Name	BAG3
Gene Alias	BAG-3, BIS, CAIR-1, MGC104307
Gene Description	BCL2-associated athanogene 3
Omim ID	603883
Gene Ontology	Hyperlink
Gene Summary	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq]
Other Designations	BAG-family molecular chaperone regulator-3 BCL2-binding athanogene 3 Bcl-2-binding protein OTTHUMP00000020599 docking protein CAIR-1

Disease

- [Alzheimer Disease](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Neoplasm Recurrence](#)

- [Neoplasms](#)
- [Psychiatric Status Rating Scales](#)
- [Schizophrenia](#)